

MICLOSULF COMBI

50% SULPHUR (S) + 0.8% MYCLOBUTANIL

WG

NON ADR
PRODUCT



ADVANTAGE

- ✓ Wettable powder easily dispersed in water.
- ✓ High compatibility in mixtures with other phytosanitary products.
- ✓ It has a synergistic effect between the two active substances.
- ✓ Reduce resistance levels.
- ✓ Fungicide with preventive, curative and eradicating action against oidium, spotting and oidiopsis.

DESCRIPTION

Product formulated with sulfur and myclobutanil in the form of wettable powder, indicated as a preventive and curative treatment of fungi.

The myclobutanil is a systemic product that prevents the development of the mycelium once the fungus is established on the plant in the incubation period, preventing new reinfections. The presence of sulfur prevents infection by new spores.

WHY SHOULD I USE IT

The combined use of sulfur with myclobutanil makes it possible to reduce the doses used of fungicidal phytosanitary products due to the synergy of the two active substances.



INSTRUCTIONS OF USE

- Use applied by dusting, taking precautions adequate to prevent risks to people and the environment.
- Compatible with the vast majority of phytosanitary products of the market, if in doubt carry out a small-scale test.
- Avoid treatment with oils 21 days before and 21 days after the application of the product.
- Do not apply foliar with temperatures above 30° C.



COMPOSITION

- **Sulphur (S)** **80 %**
- **Myclobutanil** **0,8 %**



PACKAGING

- **5 Kg**



RECOMMENDED DOSAGE

CROPS	PLAGUE	DOSE g/ha	SECURITY PERIOD
CUCURBITACEOUS	Oidium	200 - 350	14 days
	Black rot	250 - 300	10 - 14 days
GRAPES (WINE & TABLE)	Oidium	400	8 - 10 days